ANNALES ZOOLOGICI FENNICI

7ol. 18 • 1981

BLISHED JOINTLY BY

INISH ACADEMY OF SCIENCES CIETAS PRO FAUNA ET FLORA FENNICA

hank. 15 B, SF-00170 HELSINKI 17, FINLAND

SOCIETAS SCIENTIARUM FENNICA SOCIETAS BIOLOGICA FENNICA VANAMO

1001-21 10

Contents

snail communities in lakes of southern and western Finland	233—241
Alatalo, R. V.: Habitat selection of forest birds in the seasonal environment of Finland	103—114
Gyllenberg, G.: Ingestion and turnover of oil and petroleum hydrocarbons by two planctonic copepods in the Gulf of Finland	225—228
Gyllenberg, G.: Eudiaptomus gracilis (Copepoda, Calanoida): diel vertical migration in the field and diel oxygen consumption rhytm in the laboratory	229—232
Hippa, H. & Oksala, I.: Polymorphism and reproductive strategies of Enoplognatha ovata (Clerck) (Araneae, Theridiidae) in northern Europe	179—190
Kansanen, P. H.: Effects of heavy pollution on the zoobenthos in Lake Vanajavesi, southern Finland, with special reference to the meiozoobenthos	243—251
Kansanen, P. H. & Aho, J.: Changes in the macrozoobenthos associations of polluted Lake Vanajavesi, Southern Finland, over a period of 50 years	73—101
Karling, T. G.: Typhlorhynchus nanus Laidlaw, a kalyptorhynch turbellarian without proboscis (Platyhelminthes)	169—177
Korhonen, K.: Temperature in the nocturnal shelters of the redpoll (Acanthis flammea L.) and the Siberian tit (Parus cinctus Budd.) in winter	165—168
Kristiansson, H.: Distribution of the European hedgehog (Erinaceus europaeus L.) in Sweden and Finland	115—119
Pajunen, I.: Effect of long laboratory rearing on the hibernation pattern in the garden dormouse, Eliomys quercinus	129—132
Pajunen V. I.: Analysis of developmental rates in field populations of Arctocorisa carinata (Sahlb.) and Callicorisa producta (Reut.)	
(Hemiptera, Corixidae) with the aid of developmental time axes	191—197
Palmgren, P.: The mechanism of the extrinsic coxal muscles of spiders	203—207
Pankakoski, E.: Actinomycosis-like lesions in skulls of Finnish muskrats, Ondatra zibethica (L.)	209—212
Pulliainen, E.: A transect survey of small land carnivore and red fox populations on a subarctic fell in Finnish Forest Lapland over 13 winters	270—278
Pulliainen, E. & Iivanainen, J.: Winter nutrition of the willow grouse (Lagopus lagopus L.) in the extreme north of Finland	263—269
Räisänen, P., Timola, O. & Valtonen, T.: A new corer for sampling sand and moraine bottom meiofauna	133—137

Seppänen, R.: Differences in spotting pattern between populations of Aphantopus hyperantus (Lepidoptera, Satyridae) in southern Finland	1-36
Sjöberg, K. & Danell, K.: Food availability and utilization by ducks of a shallow brackish-water bay in the northern Bothnian Bay	253—261
Sunila, I.: Reproduction of Mytilus edulis L. (Bivalvia) in a brackish water area, the Gulf of Finland	121—128
Sunila, I.: Toxicity of copper and cadmium to Mytilus edulis L. (Bivalvia) in brackish water	213—223
Terhivuo, J.: Provisional atlas and population status of the Finnish amphibian and reptile species with reference to their ranges in northern Europe	139—164
Väisänen, R.: Is there more than on successional phase in the mycetophilid (Diptera) community feeding on a mushroom?	199—201
Werdelin, L.: The evolution of lynxes	37—71